

SKRIPSI
**HUBUNGAN *DISCHARGE PLANNING* TERINTEGRASI DENGAN
READMISSION PASIEN HALUSINASI
DI RSJD SURAKARTA**

Di Susun Guna Memperoleh Gelar Sarjana Keperawatan
Dalam Program Studi Keperawatan
Universitas Sahid Surakarta



Disusun oleh:
HIDAYAH NURHAYATI
2021122036

**PROGRAM STUDI KEPERAWATAN
FAKULTAS SAINS, TEKNOLOGI DAN KESEHATAN
UNIVERSITAS SAHID SURAKARTA
2023**

INTISARI

HUBUNGAN *DISCHARGE PLANNING* TERINTEGRASI DENGAN *READMISSION* PASIEN HALUSINASI DI RSJD SURAKARTA

Hidayah Nurhayati¹, Vitri Dyah², Fajar Alam²

Latar belakang : Halusinasi merupakan gejala yang paling banyak ditemukan pada pasien skizofrenia, mencapai 90%. Pasien skizofrenia seringkali mengalami *Readmission*. *Readmission* merupakan suatu penanda kualitas perawatan pasien di rumah sakit terhadap pelayanan yang diberikan. *Readmission* dapat dicegah dengan cara membuat *discharge planning* untuk pasien yang baik.

Tujuan : Untuk Mengetahui hubungan *discharge planning* terintegrasi dengan *readmission* pasien halusinasi di RSJD Surakarta.

Metode : Penelitian ini merupakan penelitian dengan pendekatan kuantitatif menggunakan desain penelitian observasional analitik, dengan rancangan penelitian *Cross Sectional*. Sampel penelitian adalah keluarga/penanggung jawab pasien halusinasi yang menerima *discharge planning* yaitu sebanyak 65 orang. Teknik pengambilan sampel dilakukan secara *purposive sampling*. Analisis data menggunakan menggunakan uji Gamma. Penelitian ini dilakukan bulan Juni 2023 di 12 ruang rawat inap RSJD Surakarta.

Hasil : Dari hasil penelitian didapatkan pelaksanaan *discharge planning* terintegrasi dengan kategori sangat baik 89,23%, baik 6,15% dan cukup 4,62%. Sedangkan *readmission* pasien halusinasi didapatkan interval *readmission* kategori I 12,31%, kategori II 4,62%, kategori III 23,08%, dan kategori IV 60%. Berdasarkan hasil analisis data hubungan *discharge planning* terintegrasi dengan *readmission* pasien halusinasi di RSJD Surakarta menggunakan uji Gamma di dapatkan nilai Sig 0.010 dan koefisien korelasi = 0.871 yang artinya terdapat hubungan yang signifikan dan korelasi sangat kuat.

Kesimpulan : Ada hubungan *discharge planning* terintegrasi dengan *readmission* pasien halusinasi.

Kata Kunci: *Discharge Planning Terintegrasi, Readmission, Halusinasi*

¹ Mahasiswa Program Studi Keperawatan, Universitas Sahid Surakarta

² Dosen Program Studi Keperawatan, Universitas Sahid Surakarta

ABSTRACT

THE CORRELATION BETWEEN INTEGRATED DISCHARGE PLANNING WITH READMISSION OF HALLUCINATE PATIENTS IN RSJD SURAKARTA

Hidayah Nurhayati¹, Vitri Dyah², Fajar Alam²

Background: Hallucinations are the most common symptom in schizophrenic patients, with 90%. Schizophrenic patients often experience readmission. Readmission is a marker of the quality of patient care at the hospital for the services provided. Readmission can be prevented by making discharge planning for good patients.

Objectives: To determine the correlation between integrated discharge planning and hallucinatory readmission in RSJD (mental hospital) Surakarta.

Methods: This research is included in the quantitative approach with an analytic observational research design and a cross sectional research design. The research sample was 65 families/persons of hallucination patients who received discharge planning. The sampling technique was carried out by purposive sampling. Data analysis used the Gamma test. This research was conducted in June 2023 in 12 inpatient rooms at the RSJD Surakarta.

Results: It was found that the implementation of integrated discharge planning was in the very good category with 89.23%, good with 6.15%, and sufficient with 4.62%. Meanwhile, the readmission of hallucinatory patients obtained the readmission interval for category I at 12.31%, category II at 4.62%, category III at 23.08%, and category IV at 60%. Based on the results of data analysis on the correlation between discharge planning integrated with the readmission of hallucinatory patients at Surakarta Hospital using the Gamma test, the Sig value was 0.010 and the correlation coefficient was 0.871. It means that there is a significant and very strong correlation.

Conclusion: There is a correlation between integrated discharge planning and hallucination patient readmission.

Keywords: Integrated Discharge Planning, Readmission, Hallucinations

¹ Students of the Nursing Department, Sahid Surakarta University

² Lecturer of Nursing Department, Sahid Surakarta University

